

## HERMAPHRODITISM IN *EURYCEA BISLINEATA*.<sup>1</sup>

INEZ WHIPPLE WILDER AND ELIZABETH BARRETT PEABODY.

The occurrence of hermaphroditism among anurans seems to be an accepted fact. Crew ('21) summarized all the recorded cases of abnormal sexual organs in frogs and states that there are forty such cases. To this Swingle ('22) has recently added one more, but finds that of his list of forty-one abnormalities only twenty-seven can be considered hermaphrodites, a sufficient number, however, coming from the hands of so severe a critic, to warrant the statement that hermaphroditism in anurans does occur. Cerruti ('07) and King ('10) following numerous earlier writers, have investigated the occurrence of the anomaly in toads with results which, though possibly subject to differences in interpretation, tend nevertheless to substantiate the existence of hermaphroditism in these forms.

No one has done for the urodeles the service which Crew has performed for the frogs, but from the paucity of published reports upon anomalies in urodeles this would not seem to be an arduous task. Thus Chapin ('15) in reporting a case of hermaphroditism found by her in *Spelerpes bislineatus* (*Eurycea bislineata*) cited reports of only two other cases of this anomaly in urodeles which had come to her attention, one that of La Valette St. George ('95) in *Triton taniatus*, the other that of Knappe ('86) in a young *Salamandra maculata*. Since the publication of Chapin's paper a third case has been reported by Krizenecky ('17) in *Triton cristatus*. Although the cases of La Valette St. George and of Krizenecky in *Triton* are unquestionably to be accepted as genuine, there is doubt concerning the nature of the anomaly reported by Knappe in *Salamandra*. Its interpretation as an hermaphrodite is apparently that of King ('10) who in summing up the reported cases of the occurrence of hermaphroditism in urodeles says that "Knappe ('86) noted the presence of a Bidder's organ in a young salamander." In the paper in question, however, following the enumeration of the species of Amphibia which he

<sup>1</sup> Contribution from the Department of Zoölogy of Smith College, No. 116.

had examined for the possible occurrence of a Bidder's organ, an enumeration which included seven species of *Anura* and two of *Urodela* (*Triton taeniatus* and *Salamandra maculata*) Knappe states definitely "Bald stellte sich heraus, dass es zur Bildung eines Bidder'schen Organs nur bei den echten Krötenarten kommt." And farther on after mentioning the unique appearance of one Bidder's organ, he says: "Eine solche Samenkörperbildung in Eikapseln des Bidder'schen Organs mit gleiches Bestimmtheit nachzuweisen, wie in dem eben beschriebenen Falle, ist mir bis jetzt nicht wieder gelungen, doch kann ich ähnliches für eine andere verwandte Thiergruppe, die Salamander, konstatiren. So liess eine in Schnittserien zerlegte Hodenabtheilung eines jungen, vielleicht zweijährigen *Salamandra maculata* nicht den geringsten Zweifel, dass dieselbe aus Eikapseln, ähnlich denen im Bidder'schen Organ der Kröten, besteht." It was thus obviously not Knappe's intention to state that he found a Bidder's organ in a salamander, but rather an appearance in the testis of a salamander like that of the unique Bidder's organ in a toad. In any case the interpretation of the condition described in the salamander as hermaphroditic, will depend upon the interpretation of the sexual nature of Bidder's organ itself. This is a matter which has been a bone of contention ever since the discovery of the organ in 1758 by Rösel von Rosenhof, and a number of theories have been advanced regarding its nature and significance.

In view of the almost universal agreement of modern writers as to the femaleness of Bidder's organ, Swingle's recent discussion ('21 and '22) of its nature is of great importance. In a discussion of the so-called transformation of sex in frogs, he claims that the theory is really based on a misinterpretation of the appearance of the cells in the Bidder's organ of toads. According to Swingle, the oviform-like cells of this organ do not represent the cells of an ovary, thus making the animal an hermaphrodite at this stage, but are, like the cells of similar appearance which occur in the pro-testis of the frogs, merely senescent male cells which are undergoing oviform degeneration. He adds further: "True hermaphroditism in frogs is a permanent and pathological condition, probably due to a mix-up in the genetic constitution of the individual,

and is not to be confused with the present problem which has to do with a normal but transitory embryological process."

The general opinion thus set forth by Swingle finds support also in a statement made by Crew ('21) who said: "Cytologically it has not been proved that the cells which constitute Bidder's organ are ovarian and there undoubtedly are reasons for questioning the generally accepted opinion that this organ is a rudimentary ovary."

The three cases of hermaphroditism already reported in urodeles, disregarding now Knappe's inconclusive report, differ from each other quite markedly. La Valette St. George's case in *Triton taniatus* was referred to by Cole ('96) as "the most complete case of hermaphroditism yet recorded among the Amphibia." The specimen was a male with perfectly distinct and independent paired ovaries, in addition to a pair of normal testes, but, however, without any traces of oviducts. The testes contained developing and fully developed sperms; and the ovaries, eggs in various stages of maturity.

The case reported by Křizenecky in *Triton cristatus* showed the presence of ova within both peripheral and internal lobules of otherwise normal testes.

The case reported by Chapin in *Eurycea bislineata* was that of an advanced larva in which the gonad was essentially male with female elements. Macroscopically, the anterior part of the left gonad, which was much reduced in size, resembled the normal testis in texture, though not in shape, while the posterior region was distinctly like an ovary. The right gonad, which was somewhat smaller than the normal testis of an individual of the same size, showed another sort of hermaphroditism. Two ova were found in the otherwise apparently normal testis, each one completely filling one lobule, which would normally contain a large number of male cells. This case showed, therefore, two ways in which female elements may be disposed in otherwise distinctly male gonads; one in the form of growing ova among the cysts of spermatogonia, and the other by a modification of a part of the gonad into a region resembling an ovary. The numerous cases of hermaphroditism which we have found in this same species are all of the same general character as that described by Chapin.

In the light of a recent article by Jordan ('22), it may be well to define our use of the term, hermaphroditism. According to Jordan, true anatomic hermaphroditism occurs "where ovary and testis are present in the same individual." Jordan regards the presence of an ovo-testis as a modification of true hermaphroditism, a condition which he designates as a type of false hermaphro-




FIG. 1. Camera lucida drawings of the ventral view of the gonads of (A) an adult male; (B) an adult hermaphrodite; and (C) an adult female ( $\times 9$ ). *Fa*, fat bodies; *m*, Müllerian duct; *ms*, mesonephros; *o*, ova (primary oöcytes). *tl*, testicular lobules; *tp*, testicular pigmentation (the two latter present but not shown in (A)); *w*, Wolffian duct. The line 1-2 shows the level of the section drawn in Fig. 4.

ditism. Were this distinction to be accepted, the term, true hermaphroditism, could be used only when referring to such a case as that of La Valette St. George's in *Triton teniatus*. There seems to be no justification, however, for this distinction of Jordan's, inasmuch as a distinct testis and ovary is but a further step in the separation of the male and female elements which, in some individuals, are still intermingled to a greater or less extent in the ovo-testis. An examination of the adult ovo-testis shown in

Fig. 1 *B* shows that a separation posterior to the testicular part on each side, such as, in fact, is slightly indicated in the right gonad, would transform each ovo-testis into a distinct testis and ovary. A female was found in which the ovary (Fig. 2) showed a number of separate parts or lobes, some connected with each other by




FIG. 2. Camera lucida drawing of the ovaries of an adult female showing an unusual lobed form ( $\times 7$ ). *M*, Müllerian duct; *ms*, mesonephros; *o*, ova; *ol*, detached lobe of ovary.

the mesovarium and others quite distinct, serving to illustrate the point that parts which are usually continuous may, through some unknown cause, become thus carried apart. Jordan's distinction in terminology seems, therefore, a somewhat arbitrary one, at least as applied to our species, and thus any individual which shows the presence of both male and female sex cells, even though these appear side by side in the same gonad, is regarded in this paper as a true hermaphrodite.

Naturally the ultimate criterion of hermaphroditism should be the production of functional germ cells of both sexes. Such a

criterion would obviate all possibility of the condition being given the interpretation which Swingle has given to the oviform cells which occur in Bidder's organ and in the larval testis (pro-testis) of frogs. In carrying out our investigation we had in mind as near an approach to this ideal as possible, and, having found examples of the condition which to us seemed unquestionably hermaphroditic in individuals of various stages up to transformation, we made a definite search for such cases among adult animals. As this search was rewarded by the discovery of one adult in which the hermaphroditic condition was beyond question, although the individual had not arrived at full sexual maturity, we feel confident that our interpretation of our cases as true hermaphrodites is correct and that the condition described cannot be considered as "a normal but transitory embryological process."

#### PERCENTAGE OF OCCURRENCE.

The determination of the percentage of occurrence of hermaphrodites with reference to that of males and females in *Eurycea bislineata* is based upon the examination of the gonads of 1113 individuals ranging from the typical larval to the adult stage. Wilder ('24) has shown that *Eurycea bislineata* is a form in which the period before transformation is considerably prolonged, covering from two to three years, although the structural changes leading toward metamorphosis are inaugurated many months previous to the actual transformation. The whole period from hatching to transformation is subdivided on the basis of structural changes into stages, the readily recognizable criteria of which, in living individuals, are as follows:

1. Postembryonic stage—Yolk still present, intestine not fully formed.
2. Typical larval stage—Intestine fully formed; no naso-lacrimal groove and no *os thyroideum*.
3. Premetamorphic stage—Open naso-lacrimal groove (in incipient phase); *os thyroideum* present; no vesicular glands in the skin.
4. Metamorphic stage—Glands of skin appearing as tiny acinous vesicles (in incipient phase), becoming rapidly larger and more conspicuous; absorption of larval structures and de-



velopment of eyelids and naso-labial groove (advanced phase).

The specimens used constituted representative collections made through a period of several years and had been preserved either in alcohol after fixation in Bouin's solution, or in formalin.

The method followed in sexing was first to examine the gonads *in situ* under a Bausch and Lomb binocular dissecting microscope with a strong artificial illumination. In many cases this was sufficient to diagnose the sex, but in those cases in which it was not, the gonad was removed and cleared *in toto* in glycerine for more careful study under the compound microscope. If in sexing one begins with adults and continues through the smaller and earlier stages, one comes finally to a point where it is practically impossible to be sure of the sex. Individuals of less than 27 mm., though frequently possessing readily sexed gonads, more often exhibit a developmental condition which might admit of various interpretations, since, at least without the use of cytological criteria, the small cells present might be either oögonia or spermatogonia. Bouin ('01) found that in *Rana temporaria* the first development of male and female germ-cells is identical as far as origin and general appearance are concerned. It may even be the case in *Eurycea*, as Okkelberg ('21) has shown for the brook lamprey, that the animal passes through a period of sex indifference before sex differentiation sets in. His observations "seem to warrant the conclusion that each larva of this species (*Entosphenus wilderi*) carries the potentiality of both sexes and that sex, therefore, is not irrevocably fixed at fertilization." He explains the development of sex in these gonads of "potentially either" sex by showing the presence in the gland of two kinds of germ cells, those manifesting a tendency towards rapid division (katabolic) and those showing a tendency to growth (anabolic). He says: "The former are regarded as having a male and the latter a female potentiality. The relative proportion of anabolic and katabolic cells determines whether the larva becomes a male or a female individual."

Since our problem, however, was not one dealing with a possible early transitory hermaphroditic condition, we have included in the calculation of the percentage of occurrence only those indi-

viduals which had definitely passed beyond the indeterminate sexual stage to a point where sex could be definitely diagnosed. The following summary is, therefore, based upon the study of gonads of animals of a length of 27 mm. and over, the arbitrary minimum of 27 mm. being taken as approximately representing the dividing line between those individuals in which the sex is still, if not indifferent, at least frequently difficult to determine, and those in which the sex is unquestionably established and recognizable.

## SUMMARY OF PERCENTAGE OF OCCURRENCE.

## Typical larval individuals (of 27 mm. and over in length)

Total number sexed.....	178
Number of males.....	86, or 48.3 %
Number of females.....	89, or 50.0 %
Number of hermaphrodites.....	3, or 1.68%

## Individuals in incipient phase of premetamorphosis

Total number sexed.....	333
Number of males.....	177, or 53.2 %
Number of females.....	152, or 45.7 %
Number of hermaphrodites.....	4, or 1.2 %

## Premetamorphic individuals (beyond incipient phase)

Total number sexed.....	256
Number of males.....	119, or 46.5 %
Number of females.....	134, or 52.3 %
Number of hermaphrodites.....	3, or 1.2 %

## Metamorphic individuals

Total number sexed.....	226
Number of males.....	115, or 50.8 %
Number of females.....	107, or 47.4 %
Number of hermaphrodites.....	4, or 1.8 %

## Adult individuals

Total number sexed.....	120
Number of males.....	77, or 64.0 %
Number of females.....	42, or 35.0 %
Number of hermaphrodites.....	1, or 0.83%

## All individuals exclusive of adults

Total number sexed.....	993
Number of males.....	497, or 50.05%
Number of females.....	482, or 48.53%
Number of hermaphrodites.....	14, or 1.41%



## All individuals

Total number sexed.....	1113
Number of males.....	574, or 51.57%
Number of females.....	524, or 47.08%
Number of hermaphrodites.....	15, or 1.35%

The variation in the percentage of occurrence of hermaphrodites in the different developmental stages has little significance because of the large probable error due to the small number of specimens examined. It should be noted that the nearest approximation to the general average, 1.35 per cent., occurs in the case of those stages in which there were the largest numbers examined. The extremely low percentage in the adult group is noteworthy and, though probably due to the small number of specimens used, may conceivably indicate a lower degree of viability in the case of hermaphrodites in adult life. The discrepancy in percentage of males and females in the adult group may also indicate a difference in viability, but is more likely to be due to the fact that the collections used were made mainly in the spring when the females would be more difficult to find because during the egg laying period they are under large rocks in the deeper water.

The fact of real significance is that of the existence of hermaphroditism in every developmental stage, since together with the approximate equality of the two sexes, it serves to eliminate any claim that the condition in question is, in this species, merely a transitory one.

This establishment of a fairly constant percentage of occurrence of hermaphroditism in *Eurycea bislineata* suggests the possibility that a search for the phenomenon in other urodeles might reveal a like frequency of occurrence. The urodeles offer an inviting field for such investigation since so little has been done with them in connection with the problem in contrast to the large amount of attention which has been given to the anurans.

DESCRIPTION OF REPRESENTATIVE CASES OF HERMAPHRODITISM  
IN *Eurycea bislineata*.

*Adult Stage.*

The one hermaphroditic adult found was an individual of 65 mm. in length which had the external characteristics of a female,

as shown by the presence of a spermatheca in the dorsal wall of the cloaca, and the shape of the head. The appearance of the gonads *in situ* is shown in Fig. 1 B. For comparison, a typical ovary and a typical testis from animals of approximately the same size and collected at the same time (July 3, 1915) are shown in Figs. 1 A and 1 C. The smaller size of both testicular and ovarian parts of the ovo-testis as compared with the typical male and female gonad respectively will be noted. Moreover, while the reproductive ducts (Wolffian and Müllerian respectively) of the normal male and female have approximately the form characteristic of




FIG. 3. Cross section showing (*m*) the Müllerian and (*w*) the Wolffian ducts in the adult hermaphrodite (cf. Fig. 1 B) ( $\times 400$ ).

the mature condition, the hermaphrodite showed macroscopically apparently a single slender straight duct upon each side following the lateral border of the mesonephros. Subsequent microscopic examinations of cross sections of this, however, revealed the presence of two ducts (cf. Fig. 3) the more lateral of which, the Müllerian, alone persists anterior to the mesonephros. This sexually indifferent condition of the ducts is identical with that shown by cross sections, made previous to this investigation, through the body of a 66 mm. immature adult male, the testes of which also correspond histologically almost exactly to the testicular portions of the hermaphroditic gonad and were thus used as a typical male control in the microscopic study.

The fat bodies in the hermaphrodite were especially large, and when the body cavity was first opened completely obscured every-

thing beneath them, making their removal necessary for the study of the gonads. The right gonad is the larger and in a macroscopic examination seems to be primarily a testis with characteristic pigmentation and conspicuous lobules. The pigmentation is, however, somewhat lighter in color than that usually found in the adult testis. The length of the testicular portion of the right gonad is 4 mm., while that of the testis shown in Fig. 1 *A* is 5.25 mm. Posterior to this testicular portion, occurs a more slender unpigmented structure in which ten large unmistakable ova, together with smaller ones, may be seen. Its general resemblance to an ovary is seen by comparison with the ovaries of the 60 mm. adult female shown in Fig. 1 *C*.

The left gonad is longer and more slender as a whole than the right. This is due to the greater length of the ovarian part, the testicular region being smaller than that of the right gonad (2.75 mm. as compared with 4 mm.). Moreover the testicular pigmentation is confined to the anterior region of the gonad and is still lighter in color than that of the right gonad. The characteristic lobules are present, but there is less differentiation of the testicular from the ovarian region, the two seeming to grade into each other insensibly. In this gonad 14 large ova are in evidence as well as numerous smaller ones. At the extreme posterior end of the left gonad there is a small semi-detached ovarian lobe.

The hermaphrodite had not been preserved originally for histological study, since the animal had been killed in 5 per cent. formalin and had been kept in this fluid since 1915. Nevertheless the gonads were sectioned, and, in spite of the excessive shrinking which is especially evident in the separation of the cysts which make up the testicular lobules, the characteristic structure of both the male and the female components was shown with unmistakable clearness.

Figure 4 shows a cross section through a region where, in a macroscopic examination, the right gonad had the appearance of a testis and the left one the appearance of an ovary. In general this section shows the typical testicular structure of the right gonad, with lobules, each made up of a number of component cysts of male cells, arranged radially about a central collecting duct. A single large ovum appears in the section, however, completely fill-

ing one of the lobules and thus apparently the equivalent of many cysts. The characteristic anabolic and katabolic nature of the female and male cells respectively is thus well exemplified. The female cell grows, the male cell divides. In the whole series of sec-




FIG. 4. Cross section through the gonads of the adult hermaphrodite at the level indicated in Fig. 1 *B* by the line 1-2 ( $\times 87$ ). The two gonads are brought nearer together in the drawing than their actual position. *Cc*, central cavity of ovarian region; *cd*, collecting duct of testicular region; *f*, follicles of both ova and testicular cysts; *mt*, spermatogonial cells in mitosis; *o*, ova (primary oöcytes); *p*, peritoneal investment; *pr*, primordial germ cells; *tc*, testicular cyst; *tl*, testicular lobule.

tions through the gonads no fewer than ten such ova were found in the testicular portion of the right gonad and six in that of the left. All were, like the one shown in Fig. 4, in an apparently normal state of development, manifesting no incipient signs of degeneration such as Crew ('21) reports to be the case in *Anura* whenever

female elements are found in parts which are primarily male in character. In fact no difference could be detected between the ova among the testicular lobules and those of the more distinctly ovarian part of the gonads except that the former had advanced further in the matter of accumulation of layers of yolk.

There was much mitotic activity in progress in the testicular lobules, the same stage of mitosis being exhibited by all the cells of a given cyst, a condition which is to be expected if one postulates their formation by repeated divisions of a single primordial spermatogonium. Thus the male elements, like the female, have every appearance of undergoing perfectly normal development. In the transition region from the testicular to the ovarian part of the gonad, small testicular lobules appear which are somewhat degenerate in character.

The posterior part of each gonad shows the typical ovarian structure as demonstrated by the section of the left gonad in Fig. 4, with large central cavity surrounded by ova, each within its layer of follicle cells.

The microscopic condition thus shown seems scarcely more advanced than that pictured and described by Chapin ('15) in the gonads of her 46 mm. hermaphroditic "larva," which, in the absence of the more exact criteria of developmental stages of the whole larval life such as we are here making use of, was designated as a "larva" in the sense that it had not yet undergone transformation. In reality it was probably an individual which was approaching metamorphosis if not in actual metamorphic condition. In general this species shows much normal variation in the developmental condition of the gonads at transformation and it is thus not surprising that one individual previous to transformation should be in the same condition as another which has already transformed.

Our adult hermaphrodite is noteworthy, not only because it shows that the condition is not merely a juvenile one, but also because so far as external characters are concerned it appears to be a female. These characters, it must be confessed, are not of a very decided nature in this species, the presence of a spermatheca being, indeed, the only unquestionable one. Moreover, the cloacal papillæ which are the characteristic male structures, might

not have appeared in so immature an individual and thus one cannot be sure that later the cloaca might not have shown male as well as female structures.

Crew ('21) in his summary of the recorded cases of abnormality of the reproductive system, says that of the 30 frogs of which sufficient details were given as to their secondary sexual characters, 25 (83.3 per cent.) were definitely and typically males; 4 others were definitely but imperfectly male (13.3 per cent.); and in the remaining case, a *Rana temporaria* described by Huxley ('20), the secondary sexual characters were female (3.3 per cent.). He says: "The abnormalities which have been recorded can be so tabulated that the first case most nearly approximates to the normal female and the last the typical male, with respect to the nature of both primary and secondary sexual characters. Thus arranged it is seen that the cases furnish an almost complete series of gradations which range from an individual almost completely female, to one almost completely male, and that the conditions found readily appear to be merely graded stages of a single process."

All of our other hermaphroditic examples of *Eurycea* were in too early a stage of development for secondary sexual characters to have appeared. However, so far as the condition of the gonads alone was concerned the same sort of graded series was found as that described by Crew in the frogs.

More thorough microscopic examination of gonads might, by disclosing occasional ova among the lobules of an otherwise normal testis or a few testicular elements concealed by the large ova of an ovary, yield a more complete seriation. At least the conditions shown by *Eurycea* indicate that in this species the hermaphroditic condition cannot be interpreted as always a modification of the same sex.

#### *Metamorphic Stage.*

In the more advanced developmental stages, as in the case of the adults, the sexing of the specimens consisted in distinguishing between a large unpigmented ovary full of bulging ova and a more slender, heavily pigmented testis with, of course, attention directed toward the detection of any combination of the two, which would mean an hermaphroditic condition. Fig. 5 *B* shows the hermaph-



roditic gonads of a 42.4 mm. animal in the advanced metamorphic stage. The gonads are essentially male so far as general shape and slight characteristic pigmentation are concerned, and there are visible in them ten unmistakable ova. Figs. 5 *A* and 5 *C* represent the gonads of a typical male and female collected on the




FIG. 5. Camera lucida drawings of the ventral view of gonads of (*A*) an incipient metamorphic male, length 40.9 mm.; (*B*) an advanced metamorphic hermaphrodite, length 42.4 mm.; and (*C*) a premetamorphic female, length 40.2 mm. ( $\times 15$ ). *Fa*, fat bodies; *ms*, mesonephros; *o*, ova; *tl*, testicular lobules; *tp*, testicular pigment; *w*, Wolffian duct. The levels of the sections of the gonads shown in Fig. 6 are indicated by the lines 1-2.

same date and of approximately the same size and stage of development, which may be used for comparison. As noted before, the smaller size of the hermaphroditic gonad, and of the ova present in it, is obvious.

Cross sections 10 micra thick were made through all three pairs of gonads and were stained, some with Delafield hæmatoxylin,






FIG. 6. Cross sections, at the levels indicated in Figure 5 by the lines 1-2, showing (A) the right gonad of an incipient metamorphic male; (B) the right gonad of an advanced metamorphic hermaphrodite; (C) the left gonad of the same individual; and (D) the right gonad of a premetamorphic female ( $\times 365$ ). *Cc*, central cavity of ovary; *cd*, collecting duct of the testis and testicular region of the hermaphroditic gonad; *d*, degenerating cells; *f*, follicles; *mt*, spermatogonial mitoses; *o*, ova (primary oöcytes); *p*, peritoneal investment; *pl*, polymorphonuclear germ cells (primary spermatogonia); *tl*, testicular lobule; *tp*, testicular pigmentation; *yo*, young primary oöcytes.

others with iron hæmatoxylin, and still others with safranin and light green. The lobules of the normal testis (Fig. 6 *A*) are made up of spermatogonia surrounded singly or in small groups with follicle cells, with which the cysts later to be formed by the division of these spermatogonia will be covered. Typical spermatogonial mitoses are seen in this and in other sections. In the hermaphroditic gonads (Figs. 6 *B* and *C*) we find a testicular structure corresponding in general to that shown by the normal testis, with typical mitoses in evidence. At the level shown in *C* in which the ovum constitutes practically half of the total diameter of the gonad, the testicular part is not quite so far advanced as in *B*, but is, in fact, in much the same condition as the more anterior region of the normal testis. The ova shown in both of these sections are typical, as will be seen by comparison with the section of the normal ovary (Fig. 6 *D*), although they are not equal in size to the larger ones of the normal ovary.

#### *Incipient Premetamorphic Stage.*

In the examination of younger stages in which little or no testicular pigment had developed, reliance for the diagnosis of the sex had to be based upon the shape of the ovary with its protruding ova to distinguish that organ from the slender testis or from the testis with female elements present in it.

Figure 7 *B* shows the general appearance of the hermaphroditic gonad of a 36 mm. incipient premetamorphic individual, and Figs. 7 *A* and *C* show gonads of a typical male and female of about the same size and developmental condition. The smaller size of the hermaphroditic gonad is again evidenced. The ova are of about the size of the smallest seen in the normal ovary.

The anterior part of the reproductive organs in each case was sectioned transversely for the purpose of studying the relation of the ducts; while the posterior part, including, in fact, the major part of the gonads themselves, was sectioned horizontally. Delafield hæmatoxylin and iron hæmatoxylin staining were used.

Histologically the developmental condition of these gonads as shown in Fig. 8 *A*, *B*, and *C* is not essentially different from that of the metamorphic stage, except that both in the normal testis and in the testicular region of the hermaphroditic gonad there are

more single spermatogonia and fewer in groups, although the lobulated structure of the gonad is evident. There were fewer instances of spermatogonial mitosis, none, in fact, discovered in the hermaphrodite, the sections of which, however, were somewhat




FIG. 7. Camera lucida drawings of the ventral view of the gonads of (A) an incipient premetamorphic male, length 37.4 mm.; (B) an incipient premetamorphic hermaphrodite, length 36 mm.; and (C) an incipient premetamorphic female, length 36.2 mm. ( $\times 15$ ). *Ms*, mesonephros, *o*, ova; *ts*, testis and testicular portion of hermaphroditic gonad.

fragmentary. In every other particular of cell arrangement and nuclear structure the testicular regions of the hermaphrodite were identical with the normal testis. The ova of the hermaphrodite,

though of smaller size, were perfectly normal in appearance. They have the typical relationship to the testicular lobules, and, owing to their earlier stage of growth, do not bulge out so conspicuously from the surface of the gonad as in the case of the metamorphic stage (cf. Fig. 6 *B* and *C*).




FIG. 8. Horizontal sections through corresponding regions of the gonads shown in Fig. 7 of (*A*) male; (*B*) hermaphrodite; (*C*) female ( $\times 356$ ). *Cc*, central cavity; *f*, follicles; *o*, ova (primary oöcytes); *p*, peritoneal investment; *pl*, polymorphonuclear germ cells (primary spermatogonia); *tl*, testicular lobule; *yo*, young primary oöcytes.

*Typical Larval Stage.*

With the younger stages, macroscopic evidence could be relied upon still less for sexing, although normal ovaries are easily recognizable if the growth period of the ova has been well entered upon, and such normal female gonads packed with growing ova of approximately uniform size have been distinguished macroscopically in larvæ as small as 25 mm. in length, though our percentage data (p. 8) did not include individuals under 27 mm. in length. The difficulty in sexing lies in the uncertainty as to the presence of male elements in gonads in which the ova are few in number but unmistakable. We have not as yet examined microscopically large numbers of gonads of young larvæ. However, in looking over our laboratory sets of serial sections of larvæ collected in September or early October, ranging in length from 17 to 25 mm. and presumably about 12 weeks old, we find that while a few of them show a condition which might be considered as sexually indifferent in that the gonads are made up of typical primordial germ cells, each with its investment of follicle cells, arranged in single rank about a central cord, a larger number of those examined, including some of the smallest individuals, show practically all of the germ cells in early maturation stages (leptonene and pachytene) or as growing oöcytes. Such an individual seems to us to be a female, since other individuals show gonads made up of more numerous, smaller germ cells grouped in such a manner as to suggest at once the incipient lobules of a typical testis. In such gonads the germ cells show no maturation phenomena, although mitosis is occasionally seen. For the most part the nuclei are either polymorphic, or in rounded form with one or more conspicuous nucleoli. As we have very few data as to the condition of the gonads the following spring, we can only express here our tentative opinion that this species, in spite of its prolonged larval life, exhibits no such early larval maturation of male germ cells synchronously with that of the female germ cells, as has been described by Swingle ('21) for *Rana catesbeiana*.

To push our power of diagnosis of sex and recognition of possible hermaphrodites back into these early stages demands as a basis, not only a careful and thorough cytological investigation of the origin and differentiation of the germ cells such as that of



Bouin ('01) and Dustin ('07) for other species of Amphibia (mainly *Anura*), and Okkelberg ('21) for the brook lamprey, but also a complete bridging over of the gap between the early developmental phenomena and the seasonal sexual phenomena of adult life.

With regard to the bearing which occasional hermaphroditism such as this has upon its regular occurrence in certain species of animals, and upon the significance of the phenomenon in general, two opposing views are held. One of these, as set forth by Doncaster ('14), regards hermaphroditism not as a primitive but as a purely secondary condition. This opinion is based mainly on the fact that the hermaphroditic species of animals are, for the most part, highly specialized ones. Sporadic hermaphroditism is thus considered an example of variation along this same direction.

The other view is that which has recently found so vigorous a supporter in Jordan ('22), that hermaphroditism, at least in the vertebrate group, is a primitive character. Jordan points out "the abundant evidence of a normal hermaphroditic condition either adult or juvenile, among lower vertebrates (*e.g.*, tunicates, cyclostomes, probably some Amphibia)," and that "the early gonads with their primordial germ cells appear identical." This view of the primitive character of hermaphroditism naturally goes hand in hand with the theory that sex determination is a matter of differential metabolism and that forms in which sex determination has become bound up in the chromosomes represent a higher stage in metabolic control of the developing organism.

Jordan points out the peculiar interest presented by the case of amphibians in this connection, since most investigators have failed to find any evidence of a sex chromosome in this group, although King ('12) describes it for *Necturus maculatus*, Levy ('15) for *Rana esculenta*, and Swingle ('17) for *Rana pipiens*. In a later paper, however, Swingle ('21) questions the correctness of his own earlier identification of an accessory chromosome in *Rana pipiens* and suggests the strong probability that Levy may also have been mistaken.

Jordan makes the suggestion that the Amphibia may constitute a group in which the evolution of the sex chromosome as a separate element can be traced, and in which also a general ten-

dency toward juvenile hermaphroditism bridges the gap between lower vertebrates where functional hermaphroditism occurs in certain classes and higher vertebrates where the condition occurs only as an anomaly. Swingle has done much to dispel the idea of juvenile hermaphroditism in the anurans. Cases of hermaphroditism such as we have here described in *Eurycea bislineata* give every evidence of being a permanent rather than juvenile condition. On the other hand, although we have thus far found no evidence that these permanent hermaphrodites arise out of an earlier condition in which the gonads have the potentiality for both sexes and may thus be regarded as capable of producing either males, females, or hermaphrodites, we do not feel that our investigation of these early stages has been sufficiently extensive to warrant us in excluding this possibility. In the absence of evidence of a chromosomal control of sex determination in this species, and indeed in Amphibia in general, one should maintain an open mind toward other possibilities. Much further investigation of the subject is therefore obviously needed and is now in progress.

#### SUMMARY.

1. True hermaphroditism occurs in approximately constant proportions in every developmental stage of *Eurycea bislineata* from typical larval to adult.
2. The percentage of occurrence of hermaphrodites in this species, based upon the examination of 1,113 individuals, is 1.35 per cent.
3. There are now on record 18 cases of hermaphroditism in urodeles. The first is that of La Valette St. George ('95) in *Triton teniatus*, the second is that discovered by Chapin ('15) in *Eurycea bislineata*, the third is that of Křizenecky ('17) in *Triton cristatus*, and the other 15 cases, in *Eurycea bislineata*, are presented in this paper.

DEPARTMENT OF ZOOLOGY, SMITH COLLEGE,  
NORTHAMPTON, MASS.  
November 13, 1923.

#### BIBLIOGRAPHY.

- Bouin, M.,  
'01 Histogenese de la Gland Genitale Femelle chez *Rana temporaria* (L.).  
Archives de Biologie, Vol. 17.



- Cerruti, A.,**  
'07 Sopra due casi di anomalia dell' apparato riproduttore nel *Bufo vulg.* Anat. Anz., Bd. XXX.
- Chapin, L. C.,**  
'15 A Case of Hermaphroditism in *Spelerpes bislineatus*. Biol. Bull., Vol. XXIX., No. 2.
- Cole, F.,**  
'96 A Case of Hermaphroditism in *Rana temporaria*. Anat. Anz., Bd. XI.
- Crew, F. A. E.,**  
'21 Sex-Reversal in Frogs and Toads. A Review of the Recorded Cases of Abnormality of the Reproductive System and an Account of a Breeding Experiment. Journ. of Genetics, Vol. 11.
- Doncaster, L.,**  
'14 The Determination of Sex. G. P. Putnam's Sons.
- Dustin, A. P.,**  
'07 Recherches sur l'origine des Gonocytes chez les Amphibiens. Archives de Biologie, Vol. 23, 1907-1908.
- Jordan, H. E.,**  
'22 The Histology of a Testis from a Case of Human Hermaphroditism, with a Consideration of the Significance of Hermaphroditism in Relation to the Question of Sex Differentiation. Amer. Journ. of Anat., Vol. 31, No. 1.
- King, H. D.,**  
'10 Some Anomalies in the Genital Organs of Toads. Amer. Journ. of Anat., Vol. 10.  
'12 Dimorphism in the Spermatozoa of *Necturus maculosa*. Anat. Rec., Vol. 6.
- Knappe, E.,**  
'86 Das Bidder'sche Organ. Morph. Jahrb., Bd. XI.
- Křizenecky, J.,**  
'17 Ein Fall von Hermaphroditismus bei *Triton cristatus*. Arch. f. Entw. Mech., Bd. 42, 4.
- Levy, F.,**  
'15 Ueber die Chromativerhältnisse in der Spermatocytogenese von *Rana esculenta*. Arch. f. mikr. Anat., Bd. 86.
- Okkelberg, P.,**  
'21 The Early History of the Germ Cells in the Brook Lamprey (*Entosphenus wilderi*) up to and including the Period of Sex Differentiation. Journ. of Morph., Vol. 35.
- St. George, la Vallette,**  
'95 Zwitterbildung beim kleinen Wassermolch (*Triton taniatus*). Arch. f. mikr. Anat., Bd. XLV.
- Swingle, W. W.,**  
'17 The Accessory Chromosome in a Frog Possessing Marked Hermaphroditic Tendencies. Biol. Bull., Vol. 33.  
'20 Neoteny and the Sexual Problem. Am. Nat., Vol. 54.  
'21 The Germ Cells of Anurans, I. The Male Sexual Cycle of *Rana catesbeiana* Larvæ. Jour. Exp. Zoöl., Vol. 32.  
'22 Is There a Transformation of Sex in Frogs? Am. Nat., Vol. LVI.
- Wildier, I. W.,**  
'24 The Relation of Growth to Metamorphosis in *Eurycea bislineata*. Journ. Exp. Zoöl., Vol. XL.